AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application.

LISTING OF CLAIMS

 (currently amended) A method of treating a fluorocompound-containing gas stream, the method comprising:

generating a plasma stream from a plasma source gas, injecting the generated-plasma stream through an aperture into a chamber,:

conveying to the plasma stream a source of OH⁻ and/or H*-ions for impinging uponcontacting the plasma stream to form heated OH⁻ and/or H* ions, comprising ions selected from the group consisting of OH⁻ and H⁺; and conveying the gas stream to the heated ions.

- 2. (currently amended) <u>TheA</u> method according to <u>Cclaim 1</u>, wherein the plasma source gas comprises an inert <u>ionisable ionizable gas</u>, for example, one of nitrogen and argon.
- 3. (currently amended) <u>TheA</u> method according to <u>Cclaim 1 or Claim 2</u>, wherein the step of generating the plasma stream from a plasma source gas further <u>comprises</u> the plasma stream is formed by generating an electric field between two electrodes, and conveying the plasma source gas between the electrodes to form the plasma stream.
- 4. (currently amended) <u>The</u>A method according to <u>Cclaim 3</u>, wherein one of the electrodes <u>provides-forms</u> at least <u>a part of a wall of said-the chamber.</u>
- 5. (currently amended) <u>TheA</u> method according to <u>Cclaim 3-or Claim 4</u>, wherein the <u>step of injecting the plasma stream into the chamber further comprises</u>

<u>injecting the</u> plasma stream is <u>injected</u> into the chamber through an aperture formed in one of the electrodes.

- 6. (currently amended) <u>TheA</u> method according to <u>any preceding-claim_1</u>, wherein <u>the step of conveying to the plasma stream a source of ions the ion source is conveyed to the plasma stream occurs prior to the injection-step of <u>injecting of the plasma stream through an aperture into the chamber.</u></u>
- 7. (currently amended) <u>The</u>A method according to <u>Cc</u>laim 6_τ wherein the <u>step of conveying ion source is conveyed</u> to the plasma stream the source of ions <u>further comprises conveying the source of ions</u> in a stream comprising the plasma source gas.
- 8. (currently amended) <u>TheA</u> method according to <u>Cclaim 6</u>, wherein the <u>source</u> <u>of ions source</u> is conveyed to the plasma stream separately from the plasma source gas.
- 9. (currently amended) <u>TheA</u> method according to <u>any of Cclaims 1 to 5</u>, wherein the <u>source of ions source</u> is conveyed to the chamber.
- 10. (currently amended) <u>The</u>A method according to <u>Cclaim 9</u>, wherein the <u>source</u> <u>of ions</u> source is conveyed into the chamber separately from the gas stream.
- 11. (currently amended) <u>The</u>A method according to <u>any preceding</u> claim <u>1</u>, wherein the gas stream is conveyed directly to the chamber for reacting with the heated ions therein.
- 12. (currently amended) <u>The</u>A method according to <u>any preceding claim 1</u>, wherein the gas stream is conveyed to the chamber separately from the plasma stream.

- 13. (currently amended) <u>TheA</u> method according to <u>any of Cclaims 1 to 6</u>, wherein the gas stream is conveyed to the heated ions through the plasma stream.
- 14. (currently amended) <u>TheA</u> method according to <u>Cclaim 13</u>, wherein the gas stream is conveyed to the plasma stream for injection into the chamber therewith.
- 15. (currently amended) A method of treating a fluorocompound-containing gas stream, the method comprising:

generating a plasma stream from a plasma source gas, adding the gas stream to the plasma stream;

injecting the plasma stream and gas stream through an aperture into a reaction-chamber, and

conveying to the plasma stream a source of OH⁻ and/or H⁺ ions comprising ions selected from the group consisting of OH⁻ and H⁺.

- 16. (currently amended) <u>The</u>A method according to <u>Cclaim 15</u>, wherein the plasma source gas comprises an inert <u>ionisable ionizable gas</u>, for example, one of nitrogen and argon.
- 17. (currently amended) <u>TheA</u> method according to <u>Cclaim 15 or Claim 16</u>, wherein the step of generating the plasma stream from the plasma source gas <u>further comprises</u> the plasma stream is formed by generating an electric field between two electrodes, and conveying the plasma source gas between the electrodes to form the plasma stream.
- 18. (currently amended) <u>The</u>A method according to <u>Calaim 17</u>, wherein one of the electrodes <u>provides forms</u> at least <u>a part of a wall of said the chamber.</u>
- 19. (currently amended) <u>TheA</u> method according to <u>Cclaim 17-or Claim 18</u>, wherein the <u>step of injecting the plasma stream and gas stream into the</u>

- <u>chamber further comprises injecting the plasma stream is injected into the chamber through an aperture formed in one of the electrodes.</u>
- 20. (currently amended) <u>TheA</u> method according to <u>any of Cclaims 15 to 19</u>, wherein the <u>step of conveying ion source is conveyed</u> to the plasma stream <u>the source of ions occurs prior</u> to the <u>step of injection injecting of the plasma stream and gas stream into the chamber.</u>
- 21. (currently amended) <u>TheA</u> method according to <u>Cclaim 20</u>, wherein the <u>step</u> of conveying ion source is conveyed to the plasma stream the source of ions <u>further comprises conveying the source of ions</u> in a gas stream comprising the plasma source gas.
- 22. (currently amended) <u>TheA</u> method according to <u>Cclaim 20</u>, wherein the ion source <u>of ions</u> is conveyed to the plasma stream separately from the plasma source gas.
- 23. (currently amended) <u>TheA</u> method according to <u>any of Cc</u>laims 15 to 20, wherein the <u>ion</u>-source <u>of ions</u> is conveyed to the plasma stream injected into the chamber.
- 24. (currently amended) <u>TheA</u> method according to <u>any of Cclaims 1514 to 20</u>, wherein the <u>ion-source of ions</u> is conveyed to the plasma stream within the gas stream.
- 25. (currently amended) <u>TheA</u> method according to <u>any preceding-claim 15</u>, wherein the plasma stream is generated <u>around or belowat atmospheric pressure</u>.
- 26. (currently amended) <u>The</u>A method according to <u>any preceding-claim 15</u>, wherein the plasma stream is generated using a dc plasma torch.

- 27. (currently amended) <u>The</u>A method according to <u>any preceding claim 15</u>, wherein the <u>ion-source</u> of ions comprises <u>one of water and water vapour</u>.
- 28. (currently amended) <u>TheA</u> method according to <u>any of Cclaims 151 to 26</u>, wherein the <u>ion-source of ions</u> comprises an alcohol, <u>for example one selected from the group consisting of methanol</u>, ethanol, propanol, propan-2-ol and butanol.
- 29. (currently amended) <u>TheA</u> method according to <u>any of Cclaims 151 to 26</u>, wherein the <u>ion-source of ions comprises a hydrogen-containing compound selected from the group consisting one-of hydrogen gas</u>, a hydrocarbon, ammonia, and a paraffin.
- 30. (currently amended) <u>The</u>A method according to <u>any preceding claim 15</u>, wherein the chamber is at a temperature in the range from ambient to 1200°C.
- 31. (currently amended) <u>TheA</u> method according to <u>any preceding-claim_15</u>, wherein the chamber is at ambient temperature.
- 32. (currently amended) <u>The</u>A method according to <u>any of Cclaims 15 to 30</u>, wherein the chamber is at a temperature in the range from 400°C to 1000°C.
- 33. (currently amended) <u>TheA</u> method according to <u>any preceding</u> claim <u>15</u>, wherein the chamber is at a pressure in the range from 10⁻³ mbar to 2000 mbar.
- 34. (currently amended) <u>TheA</u> method according to <u>any preceding</u> claim <u>15</u>, wherein the <u>step of conveying ion source is conveyed</u> into the chamber <u>the source of ions further comprises conveying the source of ions over a catalyst.</u>

- 35. (currently amended) <u>TheA</u> method according to <u>Cclaim 34</u>, wherein the catalyst comprises <u>a metal selected from the group consisting one of tungsten, silicon, iron, rhodium and platinum.</u>
- 36. (currently amended) <u>TheA</u> method according to <u>any preceding-claim 15</u>, wherein the gas stream is subsequently conveyed further comprising the step of conveying the gas stream from the chamber to a wet scrubber.
- 37. (currently amended) <u>TheA</u> method according to <u>any preceding claim 15</u>, wherein-further comprising the step of conveying the gas stream is <u>subsequently conveyed</u> from the chamber to a reactive media.
- 38. (currently amended) <u>TheA</u> method according to <u>any preceding claim 15</u>, wherein the <u>perfluorocompound containing gas stream comprises a</u> <u>perfluorocompound selected from the group consisting comprises one of CF₄, C₂F₆, CHF₃, C₃F₈, C₄F₈, NF₃ and SF₆.</u>
- 39. (currently amended) <u>An Aapparatus</u> for treating a fluorocompound-containing gas stream, the apparatus comprising:

a reaction chamber,;

means for generating a plasma stream from a plasma source gas and injecting the generated-plasma stream through an aperture into the chamber,:

means for conveying to the plasma stream a source of OH⁻ and/or H⁺ ions for impinging uponcontacting the plasma stream to form heated OH⁻ and/or H⁺-ions, comprising ions selected from the group consisting of OH⁻ and H⁺; and

means for conveying the gas stream to the heated ions.

40. (currently amended) <u>The Aapparatus according to Cclaim 39</u>, wherein the means for generating a plasma stream comprises means for generating an electric field between two electrodes, and means for conveying the plasma source gas between the electrodes to form the plasma stream.

- 41. (currently amended) <u>The Aapparatus according to Cclaim 40</u>, wherein one of the electrodes <u>provides forms</u> at least <u>a part of a wall of said the chamber.</u>
- 42. (currently amended) <u>The Aapparatus according to Cclaim 40-or Claim 41</u>, wherein the aperture is <u>located-formed</u> in one of the electrodes.
- 43. (currently amended) <u>The Aapparatus according to any of Cclaim 39 to 42</u>, wherein the means for conveying the source of OH-and/or H*-ions is arranged to convey the source of OH-and/or H* ions to the plasma stream prior to the injection of the plasma stream into the chamber.
- 44. (currently amended) <u>The Aapparatus according to any of Cclaims 39 to 42,</u> wherein the means for conveying the source of OH-and/or H* ions is arranged to convey the source of OH-and/or H* ions to the chamber.
- 45. (currently amended) The Aapparatus according to any of Cclaims 39 to 44, wherein the means for conveying the source of OH and/or H ions to the plasma stream is separate from the means for conveying the gas stream to the heated OH ons.
- 46. (currently amended) <u>The Aapparatus according to any of Cclaims 39 to 45</u>, wherein the means for conveying the gas stream to the heated OH and/or H ions is arranged to convey the gas stream directly to the chamber.
- 47. (currently amended) The Aapparatus according to any of cClaims 39 to 46, wherein the means for conveying the gas stream to the heated OH and/or H ions is arranged to convey the gas stream to the chamber through the aperture with the plasma stream.
- 48. (currently amended) <u>An Aapparatus</u> for treating a fluorocompound-containing gas stream, the apparatus comprising:

a reaction chamber,;
means for generating a plasma stream from a plasma source gas,;

International Application No. PCT/GB2005/000619

means for conveying the gas stream to the plasma stream, means for injecting the plasma stream and gas stream through an aperture into the reaction chamber, and

means for conveying to the plasma stream a source of OH⁻-and/or H⁺ ions comprising ions selected from the group consisting of OH⁻ and H⁺.

- 49. (new) The method according to claim 2 wherein the inert ionizable gas is selected from the group consisting of nitrogen and argon.
- 50. (new) The method according to claim 15 wherein the plasma stream is generated at a pressure below atmospheric pressure.
- 51. (new) The method according to claim 1 wherein the plasma stream is generated at atmospheric pressure.
- 52. (new) The method according to claim 1 wherein the plasma stream is generated at a pressure below atmospheric pressure.
- 53. (new) The method according to claim 1 wherein the plasma stream is generated using a dc plasma torch.
- 54. (new) The method according to claim 1 wherein the source of ions comprises water.
- 55. (new) The method according to claim 1 wherein the source of ions comprises an alcohol selected from the group consisting of methanol, ethanol, propanol, propan-2-ol and butanol.
- 56. (new) The method according to claim 1 wherein the source of ions comprises a hydrogen-containing compound selected from the group consisting of hydrogen gas, a hydrocarbon, ammonia, and a paraffin.

International Application No. **PCT/GB2005/000619**

- 57. (new) The method according to claim 1 wherein the chamber is at a temperature in the range from ambient to 1200°C.
- 58. (new) The method according to claim 1 wherein the chamber is at ambient temperature.
- 59. (new) The method according to claim 1 wherein the chamber is at a temperature in the range from 400°C to 1000°C.
- 60. (new) The method according to claim 1 wherein the chamber is at a pressure in the range from 10⁻³ mbar to 2000 mbar.
- 61. (new) The method according to claim 1 wherein the step of conveying into the chamber the source of ions further comprises conveying the source of ions over a catalyst.
- 62. (new) The method according to claim 61 wherein the catalyst comprises a metal selected from the group consisting of tungsten, silicon, iron, rhodium and platinum.
- 63. (new) The method according to claim 1 further comprising the step of conveying the gas stream from the chamber to a wet scrubber.
- 64. (new) The method according to claim 1 further comprising the step of conveying the gas stream from the chamber to a reactive media.
- 65. (new) The method according to claim 1 wherein the fluorocompound containing gas stream comprises a perfluorocompound selected from the group consisting of CF₄, C₂F₆, CHF₃, C₃F₈, C₄F₈, NF₃ and SF₆.